One-way MANCOVA in SPSS Statistics

Introduction

The one-way multivariate analysis of covariance (MANCOVA) can be thought of as an extension of the one-way MANOVA to incorporate a covariate or an extension of the one-way ANCOVA to incorporate multiple dependent variables. This covariate is linearly related to the dependent variables and its inclusion in the analysis removes the error variance from the variance that is shared between groups of an independent variable. A one-way MANCOVA is used to determine whether there are any statistically significant differences between the adjusted means of three or more independent groups on two or more dependent variables.

For example, you could use a one-way MANCOVA to determine whether a number of different exam performance (dependent) scores in an early levels emerged students, whilst controlling for revision time (i.e., your dependent variables would be "humanities exam performance", "science exam performance" and "mathematics exam performance", all measured from 0-100, your independent variable would be "test anxiety level", which has three groups — "low-stressed students", "moderately-stressed students" and "highly-stressed students" — and your covariate would be "revision time", measured in hours). You could run a one-way MANCOVA to determine whether that the effect of test anxiety levels on overall exam performance still exists when you have controlled for the variance explained by revision time.

The one-way MANCOVA is very useful, but it is important to realise that the one-way MANCOVA is an omnibus test statistic. It will tell you whether the groups of the independent variable statistically significantly differed based on the combined dependent variables, after controlling for the covariate, but it will not tell you how the groups differed from each other. To establish how the groups differed, you will need to perform follow-up tests to the one-way MANCOVA. For example, you could perform a one-way MANCOVA to establish that there is a difference in test anxiety levels on the combined scores from the three exams (i.e., the humanities, science and mathematics exams). However, it will not indicate whether the statistically significant difference is between the low-stressed and moderately-stressed students, between the moderately-stressed and highly-stressed students, or between the low-stressed and highly-stressed students. You will need to perform follow-up tests to the one-way MANCOVA to determine between which groups the statistically significant difference lies. **Learn how to** carry out these follow-up tests in this guide.

In this "quick start" guide, we show you how to carry out a one-way MANCOVA using SPSS Statistics, as well as interpret and report the results from this test. However, before we introduce you to this procedure, you need to understand the different assumptions that your data must meet in order for a one-way MANCOVA to give you a valid result. We discuss these assumptions next.

SPSS Statistics

Assumptions

When you choose to analyse your data using a one-way MANCOVA, a critical part of the process involves checking to make sure that the data you want to analyse can actually be analysed using a one-way MANCOVA. You need to do this because it is only appropriate to use a one-way MANCOVA if your data "passes" 9 assumptions that are required for a one-way MANCOVA to give you a valid result. Do not be surprised or disappointed if your data fails these assumptions, as this is a common occurrence when working with real-world data rather than textbook examples. However, when your data fails certain assumptions, there is often a solution to overcome this.

In practice, checking for these 9 assumptions is likely to be the most time consuming part of your analysis, requiring you to work through additional procedures in SPSS Statistics when performing your analysis, as well as spending some thinking about your data and what to do if your data violates different assumptions. In the sections below, these 9 assumptions are briefly set out:

- Assumption #1: Your two or more dependent variables should be measured at the interval or ratio level (i.e., they are continuous variables). Examples of continuous variables include revision time (measured in hours), intelligence (measured using IQ score), exam performance (measured from 0 to 100), weight (measured in kg), and so forth. You can learn more about interval and ratio variables in our article: Types of Variable.
- Assumption #2: Your two independent variables should consist of two or more categorical, independent groups (i.e., you have a nominal variable or an ordinal variable). Examples of nominal variables include gender (e.g., 2 groups: male and female), ethnicity (e.g., 3 groups: Caucasian, African American and Hispanic), profession (e.g., 5 groups: surgeon, doctor, nurse, dentist, therapist). Examples of ordinal variables include *Likert* scales (e.g., a 5-point scale with responses ranging from "strongly agree" to "strongly disagree"), where you have 5 or more categories, or even IQ scores (e.g., 101-110, 111-120, 121-130, and so forth).
- Assumption #3: You must have independent observations and a sufficient sample size. Independence of observations, or more specifically, independence of errors, is a complex statistical concept, but essentially it means that the error (or residual) in your one-way MANCOVA model should be uncorrelated with each other. For example, if you were measuring participants' intelligence, you would violate the assumption of independence of observations if you had participants in your study who were related to each other (e.g., siblings), or who knew each other, because their intelligence scores may be correlated. Fortunately, independence of observations can usually be identified from the study design and how the data was collected. Sample size and sufficiency is a complex issue, but you can find out more about it in our guide: Sample Size Calculator.
- Assumption #4: You should have no univariate or multivariate outliers. Outliers are data points within your data that do not follow the usual pattern (e.g., a participant in our example that has an unusual combination of scores on the dependent variables). Outliers can have a dramatic effect on the results of a one-way MANCOVA, so it is important to identify them. Fortunately, SPSS Statistics provides a straightforward method for identifying outliers, which we show you how to do in our guide: Identifying outliers in SPSS Statistics.
- Assumption #5: You must have multivariate normality. Unfortunately, multivariate normality is a particularly tricky assumption to test for and cannot be directly tested in SPSS Statistics. Instead, normality of each dependent variable within each group of the independent variable is often used to fix pieces as a best 'guess' as to whether there is multivariate normality. You can test for this using the **Shapiro-Wilk test of normality** in SPSS Statistics.

You can check assumptions #1, #2, #3, #7, #10, #11, #12 and #13 using SPSS Statistics. Before doing this, you should make sure that your data meets assumption #4, which is that you have no univariate or multivariate outliers. You can check assumption #4 using SPSS Statistics, as well as a statistical tests or basic assumptions correctly, the results you get when running the one-way MANCOVA might not be valid.

In the section, Test Procedure in SPSS Statistics, we illustrate the SPSS Statistics procedure to perform a one-way MANCOVA assuming that no assumptions have been violated. First, we set out the example we use to explain the one-way MANCOVA procedure in SPSS Statistics.

SPSS Statistics

Example and Data Setup in SPSS Statistics

A researcher wanted to determine whether cardiovascular health was better for normal weight individuals with higher levels of physical activity, after controlling for age, as measured by oxygen consumption (*VO2max*), and heart rate recovery (*HRR*). Participants were assigned to one of three groups: a "low physical activity" group, a "moderate physical activity" group and a "high physical activity" group. Each group contained 30 participants. Therefore, there were 90 participants in each group. In order to measure cardiovascular health, the researcher tested how body weight affected the combined dependent variables of *VO2max* and *HRR*. The data is shown in the table below (*Note: Because of heart disease, measured in mg/L) and spoken about previous (i.e., the 160 in *HERS*, measured in mg/mL).*

It was expected that increased levels of physical activity would have an overall beneficial effect on cardiovascular health, as measured by cholesterol concentration. A Machine Printer and spoints blood pressure. However the researcher knows that body weight affects cardiovascular health. As such, the researcher wanted to control for differences in the amount of body weight in the groups.

A one-way MANCOVA was carried out with cholesterol concentration, *VO2max*, *Machine Printer*, *HRR*, and systolic blood pressure, *BP*, as the dependent variables, body weight as the covariate, *Weight*, and the three physical activity groups – "low", "moderate" and "high" – as the groups of the independent variable, *Group*.

In SPSS Statistics, we create five variables: (a) the three continuous dependent variables, *VO2max*, *BP*, *HRR* and *BP*; (b) the categorical independent variable, *Group*, which has three groups: "low", "moderate" and "high"; and the covariate, *Weight*, which represents body weight. These variables have to be correctly set up in the **Variable View** and **Data View** windows of SPSS Statistics before you can carry out the one-way MANCOVA.

Note: You will also need to create one additional variable, *id*, to act as a case number. This variable is required to test whether there are any multivariate outliers (i.e., part of Assumption #4 above).

SPSS Statistics

Test Procedure in SPSS Statistics

The five steps below show you how to analyse your data using a one-way MANCOVA in SPSS Statistics when the 11 assumptions in the previous section, Assumptions, have not been violated. At the end of these five steps, we show you how to interpret the results from this test.

Note: Since some of the options in the **General Linear Model** > **Multivariate...** procedure changed in SPSS Statistics **version 25**, we include the instructions for **version 24** and **earlier versions** of SPSS Statistics as well. Therefore, any differences in screenshots for **version 24** and **earlier versions** in the six steps below are shown in the yellow notes like this one at the end of each step. If you are unsure which version of SPSS Statistics you are using, see our guide: Determining your version of SPSS Statistics.


Note: In **version 27** and the **subscription version**, SPSS Statistics introduced a new look to their interface called **"SPSS Light"**, replacing the previous look for **versions 26** and **earlier versions**, which was called **"SPSS Standard"**. Therefore, if you have **versions 27**, **28**, **29** or the **subscription version** of SPSS Statistics, the first step you see will look like this. However, the procedures for **versions 27**, **28**, **29** and the **subscription version** are identical.

1 Click **Analyze** > **General Linear Model** > **Multivariate...** on the main menu, as shown below:



Published with written permission from SPSS Statistics, IBM Corporation.

You will be presented with the **Multivariate** dialogue box, as shown below:



Published with written permission from SPSS Statistics, IBM Corporation.

2 Transfer the dependent variables, *VO2max*, *BP*, *HRR* and *BP*, into the **Dependent Variables:** box, the independent variable, *Group*, into the **Fixed Factor(s):** box, and the covariate variable, *Weight*, into the **Covariate(s):** box using the relevant buttons. You will end up with a screen similar to the following:



Published with written permission from SPSS Statistics, IBM Corporation.

3 If you have SPSS Statistics **versions 25, 26** or **27** (or the **subscription version** of SPSS Statistics), you can instruct SPSS Statistics to run a **post hoc test** to follow up a **statistically significant** one-way MANCOVA analysis using the **Multivariate Estimated Marginal Means** and **Multivariate Options** dialogue boxes. These can be accessed by clicking on the Options... and Options... buttons, which will launch the **Multivariate Estimated Marginal Means** and **Multivariate Options** dialogue boxes, as shown below:



Published with written permission from SPSS Statistics, IBM Corporation.



Published with written permission from SPSS Statistics, IBM Corporation.

Note: If you have SPSS Statistics **version 24** or an **earlier version** of SPSS Statistics, the options to run a **post hoc test** to follow up a **statistically significant** one-way MANCOVA analysis are accessed by clicking on the Options... button, which will launch the **Multivariate: Options** dialogue box, as shown below:



In this "quick start" guide, we simply show you how to interpret the main one-way MANCOVA result and not the follow-up procedures. Therefore, you can move onto the next step below.

4 Click on the Continue button. This will return you to the **Multivariate** dialogue box.

5 Click on the OK button to generate the results for the one-way MANCOVA.

SPSS Statistics

Interpreting the Results of the One-Way MANCOVA

SPSS Statistics produces a number of different tables in its one-way MANCOVA analysis, but you only need to interpret the **Multivariate Tests** table to determine whether your one-way MANCOVA is statistically significant. To reiterate, the analysis below assumes that your data has already met the 11 assumptions required for a one-way MANCOVA to give you a valid result.

SPSS Statistics

Multivariate Tests

The main purpose of running a one-way MANCOVA is to establish whether the groups of the independent variable, *group*, are statistically significantly different on the dependent variables (i.e., *VO2max*, *BP*, *HRR* and *BP*, collectively) after controlling for a covariate, *Weight*. The **Multivariate Tests** table in the SPSS Statistics output below lists the various multivariate statistical tests that can be used to determine whether the groups of the independent variable in terms of the **combined** dependent variable after adjusting for the continuous covariate). For example, it would suggest that the combined mean scores of our dependent variables, *VO2max*, *BP*, *HRR* and *BP*, which have been adjusted for the continuous covariate, *Weight*, differ between the three groups of our independent variable, *group*, (i.e., the "low", "Moderate" and "High" physical activity groups). Therefore, in order to interpret the results from the one-way MANCOVA, you need to look at the **Multivariate Tests** table, as highlighted below:

Multivariate Tests[a]

Effect		Value	F	Hypothesis df	Error df	Sig.	Partial Eta Squared
Intercept	Pillai's Trace	.999	24.119[b]	5.000	114.000	.000	.514
	Wilks' Lambda	.001	24.119[b]	5.000	114.000	.000	.514
	Hotelling's Trace	1.057	24.119[b]	5.000	114.000	.000	.514
	Roy's Largest Root	1.057	24.119[b]	5.000	114.000	.000	.514
weight	Pillai's Trace	.111	3.124[b]	5.000	114.000	.011	.121
	Wilks' Lambda	.111	3.124[b]	5.000	114.000	.011	.121
	Hotelling's Trace	.171	3.124[b]	5.000	114.000	.011	.121
	Roy's Largest Root	.171	3.124[b]	5.000	114.000	.011	.121
group	Pillai's Trace	.358	21.341	6.000	230.000	.000	.358
	Wilks' Lambda	.644	20.907[b]	6.000	228.000	.000	.355
	Hotelling's Trace	.537	20.707	6.000	226.000	.000	.355
	Roy's Largest Root	.517	19.812[c]	3.000	115.000	.000	.341

a. Design: Intercept + weight + group
b. Exact statistic
c. The statistic is an upper bound on F that yields a lower bound on the significance level.

Published with written permission from SPSS Statistics, IBM Corporation.

Note: We have highlighted the rows within the **group** heading. This row heading will have the same name as your independent variable, in this example, it is therefore **group** because this is the name of our independent variable, *group*. Therefore, when you analyse your own data, look for the row heading in the **Multivariate Tests** table that matches the name of your independent variable.

The different names given to each row – **Pillai's Trace**, **Wilks' Lambda**, **Hotelling's Trace** and **Roy's Largest Root** – are the names of the different multivariate statistical tests that can be used to determine whether your one-way MANCOVA is statistically significant. Which one you should use depends on a number of factors, but most researchers tend to report **Wilks' Lambda** (Λ) as it is the most commonly reported multivariate statistic and provides you with an effect size. However, you have the added complication of having to decide which multivariate *F*-statistic to use, especially because they do not always provide the same answer. The most commonly recommended multivariate statistic to use is **Wilks' Lambda** (Λ) and this is what will be used in this example.

Therefore, the result for the one-way MANCOVA is found on the "**Wilks' Lambda**" row within the "**group**" row of the **Multivariate Tests** table, as highlighted below:

Multivariate Tests[a]

Effect		Value	F	Hypothesis df	Error df	Sig.	Partial Eta Squared
Intercept	Pillai's Trace	.999	24.119[b]	5.000	114.000	.000	.514
	Wilks' Lambda	.001	24.119[b]	5.000	114.000	.000	.514
	Hotelling's Trace	1.057	24.119[b]	5.000	114.000	.000	.514
	Roy's Largest Root	1.057	24.119[b]	5.000	114.000	.000	.514
weight	Pillai's Trace	.111	3.124[b]	5.000	114.000	.011	.121
	Wilks' Lambda	.111	3.124[b]	5.000	114.000	.011	.121
	Hotelling's Trace	.171	3.124[b]	5.000	114.000	.011	.121
	Roy's Largest Root	.171	3.124[b]	5.000	114.000	.011	.121
group	Pillai's Trace	.358	21.341	6.000	230.000	.000	.358
	Wilks' Lambda	.644	20.907[b]	6.000	228.000	.000	.355
	Hotelling's Trace	.537	20.707	6.000	226.000	.000	.355
	Roy's Largest Root	.517	19.812[c]	3.000	115.000	.000	.341

a. Design: Intercept + weight + group
b. Exact statistic
c. The statistic is an upper bound on F that yields a lower bound on the significance level.

Published with written permission from SPSS Statistics, IBM Corporation.

You can see that the statistical significance is .000 (i.e., the "**Sig.**" column), which means that *p* = .000. Since .000 ≤ *α* = .050, the groups of the independent variable, *group*, (i.e., the "low", "moderate" and "high" physical activity groups) are statistically significant differences in the combined dependent variables (measured using *VO2max*, *BP*, *HRR* and *BP*) after controlling for weight.

SPSS Statistics will also report an **effect size**, called **partial eta squared** (i.e., partial η^2). This value was found in the "**Partial Eta Squared**" column of the **Multivariate Tests** table. At present, there are no agreed upon definitions of what constitutes a strong (or otherwise) effect size (Huberty & Olejnik, 2006).

SPSS Statistics

Reporting the Results of the One-Way MANCOVA

You could report a statistically significant one-way MANCOVA result as follows:

General 49%

There was a statistically significant difference between the physical activity groups on the combined dependent variables after controlling for weight, $F(5, 114) = 24.119$, $p = .000$, Wilks' $\Lambda = .001$, partial $\eta^2 = .514$.

In the statement above, the *F*(5, 114) = 24.119, *p* = .000, Wilks' Λ = .001, partial η^2 = .514 part has the following meaning:

	Column name	Column meaning
	F	Indicates that we are comparing to an *F*-distribution (*F* test).
	5 in (5, 114)	Indicates the degrees of freedom for the effect of the one-way MANCOVA.
	114 in (5, 114)	Indicates the degrees of freedom for the error term for Wilks' Lambda.
	24.047	Indicates the obtained value of the *F*-statistic (obtained *F*-value) for Wilks' Lambda.
	p = .0005	Indicates the probability of obtaining the observed *F*-value given the null hypothesis is true.
	Wilks' Λ = .259	Indicates the value of Wilks' Lambda.
	partial η^2 = .481	A measure of effect size for Wilks' Lambda.

Alternatively, if the one-way MANCOVA result **was not** statistically significant, you could report the result as follows:

General 49%

There was no statistically significant difference between the physical activity groups on the combined dependent variables after controlling for weight, $F(5, 114) = 1.279$, $p = .281$, Wilks' $\Lambda = .944$, partial $\eta^2 = .056$.

Note: The values used for the statement above were made up to provide an example.

SPSS Statistics

Following-up a Statistically Significant One-Way MANCOVA

How to follow up a statistically significant one-way MANCOVA result is controversial. One method that is recommended, which is also the default action of SPSS Statistics, is to follow up with univariate statistical tests (Pituch & Stevens, 2016). This involves two steps:

- STEP ONE: If the one-way MANCOVA is statistically significant, this suggests that there is a statistically significant adjusted mean difference between the groups of the independent variable in terms of the combined dependent variables (after adjusting for the covariates). However, you do not know whether this is due to one dependent variable or all dependent variables. Therefore, you need to investigate the adjusted mean differences between the groups of the independent variable in terms of each dependent variable. This is the first follow-up test that you need to perform.
- STEP TWO: If any of the follow-up analyses in **STEP ONE** (i.e., univariate tests) are statistically significant, you can follow this up by examining all possible pairwise comparisons between the groups of your independent variable. This will indicate which of the groups of the independent variable are statistically significantly different in terms of the adjusted group means on the dependent variable (i.e., after adjusting for the covariate). For example, these pairwise comparisons will explain whether each dependent variable, after adjusting for the covariate, is significantly different between the low physical activity group (i.e., Group 1), Group 2 and Group 3. This is the second follow-up test that you need to perform.

If the one-way MANCOVA is **not** statistically significant, you would normally not follow up the one-way MANCOVA with any further tests.